

DUKE-CGTP-06-04

hep-th/0610209

October 2006

The Landau–Ginzburg to Calabi–Yau Dictionary for D-Branes

Paul S. Aspinwall

Center for Geometry and Theoretical Physics, Box 90318
Duke University, Durham, NC 27708-0318

Abstract

Based on work by Orlov, we give a precise recipe for mapping between B-type D-branes in a Landau–Ginzburg orbifold model (or Gepner model) and the corresponding large-radius Calabi–Yau manifold. The D-branes in Landau–Ginzburg theories correspond to matrix factorizations and the D-branes on the Calabi–Yau manifolds are objects in the derived category. We give several examples including branes on quotient singularities associated to weighted projective spaces. We are able to confirm several conjectures and statements in the literature.

1 Introduction

B-type D-branes allow remarkable insight into the phase picture [1, 2] of Calabi–Yau manifolds. Given a smooth Calabi–Yau manifold X , it is known that B-type D-branes are described by the derived¹ category of coherent sheaves [3–6]. However, as one varies the Kähler form on X , one can move into other phases where the Calabi–Yau and its D-branes have another interpretation.

The most obvious “other phase” is the one studied originally in [1], namely the Landau–Ginzburg phase. Note that this phase need not exist, but we will restrict attention here to cases where it does. This Landau–Ginzburg theory also may, or may not, have a Gepner model description [7] corresponding to the construction of [8].

Certain D-branes in Gepner models were studied in [9, 10]. A more general picture of D-branes in terms of Landau–Ginzburg theories was then given in [11–14]. Suppose X is described by a hypersurface $W = 0$ in a (weighted) projective space. The corresponding Landau–Ginzburg theory then has a superpotential W . B-type D-branes in this theory correspond to matrix factorizations:

$$A.B = W.\text{id}, \tag{1}$$

where A and B are matrices (of arbitrary dimension). The correspondence between Gepner model D-branes and such matrix factorizations was described in [15].

Since the B-model is invariant to deformations of the Kähler form, the category of (topological) B-type D-branes in the Landau–Ginzburg phase must be equivalent to the category of such branes in the large radius Calabi–Yau phase. It is thus natural to ask how this correspondence works exactly. Namely, given a matrix factorization, how does one find the corresponding geometric D-brane on X ? Or, given a geometric brane, what is the corresponding matrix factorization?

There has been some progress in answering this question in several examples [15–21] but no systematic machine to convert one picture into another has been described.

Orlov [22] recently proved the equivalence of the category of matrix factorizations and the derived category of coherent sheaves on the corresponding hypersurface. Orlov’s proof is actually constructive and allows for an explicit mapping between these categories. In this paper we will demonstrate how this works.

The mathematics involved is quite technical but it seems to be very naturally tied to the phase structure of $N = (2, 2)$ theories. The principal concepts involved are quotient triangulated categories and semiorthogonal decompositions. The categories of D-branes in the Landau–Ginzburg phase and Calabi–Yau phases are both quotients of the same initial category. The semiorthogonal decompositions with respect to these quotients allows for an explicit map between the two categories.

In section 2 we will review the necessary material we require from [22] and give the recipe for turning matrix factorizations into complexes of sheaves and *vice versa*. In some cases, in particular for single points and rational curves, the process is relatively easy and we discuss

¹All derived categories in this paper will be *bounded*.

this in section 3. The Recknagel–Schomerus branes (or RS-branes) and their bound states are discussed in section 4. The case of D-branes on quotient singularities is discussed in section 5 and we give our concluding remarks in section 6.

2 Orlov’s construction

In this section we review Orlov’s construction [22] and show how it can be used to explicitly map between matrix factorizations and coherent sheaves. We refer to Orlov’s papers [22, 23] for all the proofs of the assertions below.

One proceeds through a sequence of equivalences between triangulated categories which we describe in turn. Let us begin with a graded polynomial ring

$$B = k[x_0, \dots, x_{n-1}], \quad (2)$$

where k is a field (i.e., $\mathbb{C}$ for our purposes). One may choose to give all the variables grade one, but we can also consider weighted projective spaces by allowing arbitrary degrees. One then defines the superpotential of the Landau–Ginzburg theory

$$W = f(x_0, \dots, x_{n-1}), \quad (3)$$

as a homogeneous polynomial of total degree d . We then have a quotient ring

$$A = \frac{B}{(W)}. \quad (4)$$

This defines a hypersurface X given by $W = 0$ in a (weighted) projective space. In the language of algebraic geometry, $X = \text{Proj } A$.

2.1 DGrB(W)

The category of B-type D-branes in a Landau–Ginzburg theory was described in [24]. Objects $\bar{P}$ are ordered pairs of free B -modules of arbitrary but equal rank with maps between them going in each direction:

$$\bar{P} = \left(P_1 \begin{smallmatrix} \xrightarrow{p_1} \\ \xleftarrow{p_0} \end{smallmatrix} P_0 \right). \quad (5)$$

The two maps satisfy the matrix factorization condition

$$p_0 p_1 = p_1 p_0 = W \cdot \text{id}. \quad (6)$$

A map $f : \bar{P} \rightarrow \bar{Q}$ is simply a pair of maps $f_0 : P_0 \rightarrow Q_0$ and $f_1 : P_1 \rightarrow Q_1$ such that all squares commute. Such a map is said to be null-homotopic if there are maps $s_0 : P_0 \rightarrow Q_1$ and $s_1 : P_1 \rightarrow Q_0$ such that

$$f_0 = s_1 p_0 + q_1 s_0, \quad f_1 = q_0 s_1 + s_0 p_1. \quad (7)$$

The category of D-branes is given by the homotopy category obtained by identifying morphisms with maps modulo null-homotopies. The Hilbert space of open strings in the topological B-model between two branes is given by the space of morphisms in this category. Note that if either p_0 or p_1 is the identity map, it follows from this construction that $\bar{P}$ is equivalent to 0 in this category.

For the Landau–Ginzburg–Calabi–Yau correspondence, one must orbifold the Landau–Ginzburg theory by $\mathbb{Z}_d$. It was noted in [25] that the effect of this orbifolding is to put a well-defined grading structure on the above category. That is, P_0 and P_1 are graded B -modules and one defines p_0 to have degree d and p_1 to have degree 0. Open strings are maps of degree zero.

We denote this category of D-branes on a Landau–Ginzburg orbifold $\mathrm{DGrB}(W)$. One can show that this category is a triangulated category. In particular the shift functor $[1]$ is defined by

$$\bar{P}[1] = \left(P_0 \begin{smallmatrix} \xrightarrow{p_0} \\ \xleftarrow{p_1} \end{smallmatrix} P_1(d) \right), \quad (8)$$

where (d) denotes a shift in grading. That is, if M is a graded module then $(M(d))_n = M_{d+n}$. Note in particular that

$$\bar{P}[2] = \bar{P}(d). \quad (9)$$

2.2 $\mathbf{D}_{\mathrm{Sg}}^{\mathrm{gr}}(A)$

Orlov then shows that $\mathrm{DGrB}(W)$ is equivalent to $\mathbf{D}_{\mathrm{Sg}}^{\mathrm{gr}}(A)$ introduced in [23] which is defined as follows. Let $\mathrm{gr}\text{-}A$ be the category of graded A -modules.² Note that the morphisms in $\mathrm{gr}\text{-}A$ are module homomorphisms of *degree zero*. $\mathbf{D}(\mathrm{gr}\text{-}A)$ is then the bounded derived category of graded A -modules.

Now let $\mathfrak{P}\mathrm{erf}(A)$ be the full subcategory of $\mathbf{D}(\mathrm{gr}\text{-}A)$ of objects which may be represented by finite-length complexes of free A -modules of finite rank.

One of the key ideas we require in this paper is the notion of a quotient triangulated category. Given a triangulated category $\mathbf{D}$ and full triangulated subcategory $\mathbf{N}$, we define the quotient $\mathbf{D}/\mathbf{N}$ as follows. The objects in $\mathbf{D}/\mathbf{N}$ are the same as the objects in $\mathbf{D}$. Consider the set of morphisms Σ in $\mathbf{D}$ whose mapping cones lie in $\mathbf{N}$. In other words $f : \mathbf{a} \rightarrow \mathbf{b}$ lies in Σ if and only if we have a distinguished triangle

$$\begin{array}{ccc} \mathbf{a} & \xrightarrow{f} & \mathbf{b} \\ & \searrow & \swarrow \\ & \mathbf{n} & \end{array} \quad (10)$$

where $\mathbf{n}$ is an object in $\mathbf{N}$. The morphisms in $\mathbf{D}/\mathbf{N}$ are then defined by “localizing” on the set Σ . That is, we invert the elements of Σ in the same way that quasi-isomorphisms are inverted in defining the derived category.

²As A is commutative, we need not distinguish between left and right actions.

Note, in particular, that the zero map $0 \rightarrow \mathfrak{n}$ is in Σ so that any element of $\mathbf{N}$ is isomorphic to zero in $\mathbf{D}/\mathbf{N}$.

We then define

$$\mathbf{D}_{\text{Sg}}^{\text{gr}}(A) = \frac{\mathbf{D}(\text{gr-}A)}{\mathfrak{P}\text{erf}(A)}. \quad (11)$$

The correspondence between $\text{DGrB}(W)$ and $\mathbf{D}_{\text{Sg}}^{\text{gr}}(A)$ may be seen explicitly following a result of Eisenbud [26]. Consider any A -module M and compute a minimal free resolution. Eisenbud showed that, if such a resolution has infinite length, then ultimately it is periodic with period two. That is, we have an exact sequence:

$$\begin{aligned} \cdots \xrightarrow{p_1} P_0 \xrightarrow{p_0} P_1 \xrightarrow{p_1} P_0 \xrightarrow{p_0} P_1 \xrightarrow{p_1} \cdots \\ \cdots \longrightarrow F^{-1} \longrightarrow F^0 \longrightarrow M \longrightarrow 0, \end{aligned} \quad (12)$$

where P_0 , P_1 and F^k are free A -modules, and p_0 and p_1 satisfy the matrix factorization condition (6).

So, given any A -module, we may map it to a matrix factorization by computing a minimal free resolution. Clearly if two A -modules “differ” by an A -module with finite free resolution, they will produce the same matrix factorization. Extending this to complexes gives the map from $\mathbf{D}_{\text{Sg}}^{\text{gr}}(A)$ to $\text{DGrB}(W)$. We will see examples of this later.

2.3 $\mathbf{D}(\text{qgr-}A)$

The next category we need to define is $\mathbf{D}(\text{qgr-}A)$. An A -module is said to be *torsion* if it is finite-dimensional as a vector space over k . That is, it is annihilated by x_i^N for any i and sufficiently large N . Let $\mathbf{D}(\text{qgr-}A)$ be the quotient of $\mathbf{D}(\text{gr-}A)$ by the full subcategory given by complexes of torsion modules.

It is a standard result due to Serre [27] that $\mathbf{D}(\text{qgr-}A)$ is equivalent to the derived category of coherent sheaves $\mathbf{D}(X)$.³ It is fairly easy to see why this should be so. Start with the fact that $\mathbb{P}^{n-1}$ with homogeneous coordinates $[x_0, \dots, x_{n-1}]$ is constructed by $(\mathbb{C}^n - (0, 0, \dots, 0))/\mathbb{C}^*$. A standard construction in algebraic geometry can be used to turn a module into a sheaf. We refer to [28] for details. In order to define a module on a projective variety we demand that the module has a graded structure in order to be compatible with the division by $\mathbb{C}^*$. Then note that any module producing a sheaf supported at the origin $(0, 0, \dots, 0)$ will yield a trivial sheaf in $\mathbb{P}^{n-1}$. Ignoring such sheaves amounts to taking a quotient by torsion sheaves.

So we have arrived at the statement that the equivalence between Landau–Ginzburg D-branes and large radius Calabi–Yau D-branes is an equivalence between $\mathbf{D}_{\text{Sg}}^{\text{gr}}(A)$ and $\mathbf{D}(\text{qgr-}A)$. That is, *we need to consider an equivalence between two different quotients of the derived category of graded A -modules*. This step is the only really substantial one in understanding how to map between the two different kinds of D-branes.

³See also exercise 5.9 in section II of [28]. Also one should more correctly use the language of stacks for the weighted projective space case. We discuss this more in section 5.

2.4 Semiorthogonal Decompositions

The key concept in understanding the map between $\mathbf{D}_{\text{Sg}}^{\text{gr}}(A)$ and $\mathbf{D}(\text{qgr-}A)$ is that of a *semiorthogonal decomposition*. Let $\mathcal{C}$ be a triangulated category. We say that $\mathcal{C} = \langle \mathbf{A}, \mathbf{B} \rangle$ is a semiorthogonal decomposition⁴ of $\mathcal{C}$ if the following three conditions are met:

1. $\mathbf{A}$ and $\mathbf{B}$ are full triangulated subcategories of $\mathcal{C}$.
2. For any object c in $\mathcal{C}$, there is a distinguished triangle

$$\begin{array}{ccc} \mathbf{a} & \xrightarrow{[1]} & \mathbf{b} \\ & \searrow & \swarrow \\ & \mathbf{c} & \end{array} \quad (13)$$

in $\mathcal{C}$ where $\mathbf{a}$ is an object in $\mathbf{A}$ and $\mathbf{b}$ is an object in $\mathbf{B}$.

3. $\text{Hom}_{\mathcal{C}}(\mathbf{b}, \mathbf{a}) = 0$ for any $\mathbf{a}$ in $\mathbf{A}$ and any $\mathbf{b}$ in $\mathbf{B}$.

Fix an integer i and let $\text{gr-}A_{\geq i}$ be the category of graded A -modules M such that M_j is only nonzero if $j \geq i$. Then $\mathbf{D}(\text{gr-}A_{\geq i})$ is a full subcategory of $\mathbf{D}(\text{gr-}A)$. Orlov then proved that there are two interesting semiorthogonal decompositions

$$\begin{aligned} \mathbf{D}(\text{gr-}A_{\geq i}) &= \langle \mathcal{D}_i, \mathcal{S}_{\geq i} \rangle \\ \mathbf{D}(\text{gr-}A_{\geq i}) &= \langle \mathcal{P}_{\geq i}, \mathcal{T}_i \rangle. \end{aligned} \quad (14)$$

The subcategories in this decomposition are defined as follows. Let $\mathbf{k}$ be the A -module defined as the one-dimensional vector space k in grade 0 which is annihilated by x_j for any j . Then $\mathbf{k}(-e)$ is the corresponding one-dimensional space with grade e . We define $\mathcal{S}_{\geq i}$ as the triangulated subcategory of $\mathbf{D}(\text{gr-}A_{\geq i})$ generated⁵ by $\mathbf{k}(-e)$ for $e \geq i$. As we will discuss further in section 4.2, the objects in $\mathcal{S}_{\geq i}$ correspond to bounded complexes of torsion modules in $\mathbf{D}(\text{gr-}A_{\geq i})$.

$\mathcal{P}_{\geq i}$ is defined as the triangulated subcategory of $\mathbf{D}(\text{gr-}A_{\geq i})$ generated by $A(-e)$ for $e \geq i$. Clearly objects in $\mathcal{P}_{\geq i}$ are the objects in $\mathbf{D}(\text{gr-}A_{\geq i})$ which have a finite length free resolution. The subcategories $\mathcal{D}_i$ and $\mathcal{T}_i$ are then defined from the decompositions (14).

We now claim that $\mathcal{T}_i$ is equivalent to $\mathbf{D}_{\text{Sg}}^{\text{gr}}(A)$. To see this note that the category of A -modules with finite free resolution has a semiorthogonal decomposition $\langle \mathcal{P}_{\geq i}, \mathcal{P}_{< i} \rangle$, where $\mathcal{P}_{< i}$ is generated by $A(-e)$ for $e < i$. One can then demonstrate the following equivalences:

$$\begin{aligned} \mathbf{D}_{\text{Sg}}^{\text{gr}}(A) &= \frac{\mathbf{D}(\text{gr-}A)}{\langle \mathcal{P}_{\geq i}, \mathcal{P}_{< i} \rangle} \\ &= \frac{\langle \mathbf{D}(\text{gr-}A_{\geq i}), \mathcal{P}_{< i} \rangle}{\langle \mathcal{P}_{\geq i}, \mathcal{P}_{< i} \rangle} \\ &= \frac{\mathbf{D}(\text{gr-}A_{\geq i})}{\mathcal{P}_{\geq i}} \\ &= \mathcal{T}_i \end{aligned} \quad (15)$$

⁴The semiorthogonal decompositions in this paper will all be “weak” in the sense of [22].

⁵We will always assume *finitely* generated.

Let $\pi : \mathbf{D}(\text{gr-}A) \rightarrow \mathbf{D}(\text{qgr-}A)$ be the quotient map. We now have an explicit map

$$\Phi_i : \mathbf{D}_{\text{Sg}}^{\text{gr}}(A) \xrightarrow{\sim} \mathcal{T}_i \xrightarrow{\pi} \mathbf{D}(\text{qgr-}A). \quad (16)$$

Orlov shows that if X is a Calabi–Yau manifold, then Φ_i is an equivalence and thus he proves that topological D-branes in a Landau–Ginzburg theory are the same as those on Calabi–Yau manifold.

Similarly one can show that $\mathcal{D}_i$ is isomorphic to $\mathbf{D}(\text{qgr-}A)$ and we have a map

$$\Psi_i : \mathbf{D}(\text{qgr-}A) \xrightarrow{\sim} \mathcal{D}_i \xrightarrow{\pi} \mathbf{D}_{\text{Sg}}^{\text{gr}}(A), \quad (17)$$

which takes geometric D-branes on a Calabi–Yau manifold to matrix factorizations.

2.5 The explicit map

We now have a systematic method for computing the representation of a D-brane as a complex of sheaves given the D-brane as a matrix factorization. This map is essentially given above by (15) and (16). The steps are as follows:

1. Given a matrix factorization of the form (5) we need to find any A -module M with a resolution of the form (12). This could be done by setting M equal to the cokernel of the map p_1 , although we will not use this method below.
2. Use the semiorthogonal decomposition $\mathbf{D}(\text{gr-}A) = \langle \mathbf{D}(\text{gr-}A_{\geq i}), \mathcal{P}_{< i} \rangle$ to decompose M into a part M' that lives in $\mathbf{D}(\text{gr-}A_{\geq i})$ and a part, which we discard, that lives in $\mathcal{P}_{< i}$.
3. Use the semiorthogonal decomposition $\mathbf{D}(\text{gr-}A_{\geq i}) = \langle \mathcal{P}_{\geq i}, \mathcal{T}_i \rangle$ to decompose M' into a part M'' which lies in $\mathcal{T}_i$ and a part, which we discard, that lies in $\mathcal{P}_{\geq i}$.
4. The equivalence class of M'' in $\mathbf{D}(\text{qgr-}A) = \mathbf{D}(X)$ yields the desired complex of sheaves.

Similarly we may start with a sheaf (or a complex of sheaves) on X and find the corresponding matrix factorization with the following steps:

- 1'. Given a coherent sheaf on X , we construct the corresponding graded A -module M in the standard way.
- 2'. Use the semiorthogonal decomposition $\mathbf{D}(\text{gr-}A) = \langle \mathcal{S}_{< i}, \mathbf{D}(\text{gr-}A_{\geq i}) \rangle$ to decompose M into a part M' that lives in $\mathbf{D}(\text{gr-}A_{\geq i})$ and a part, which we discard, that lives in $\mathcal{S}_{< i}$.
- 3'. Use the semiorthogonal decomposition $\mathbf{D}(\text{gr-}A_{\geq i}) = \langle \mathcal{D}_i, \mathcal{S}_{\geq i} \rangle$ to decompose M' into a part M'' which lies in $\mathcal{D}_i$ and a part, which we discard, that lies in $\mathcal{S}_{\geq i}$.
- 4'. Compute a minimal free resolution of M'' and use its asymptotic form to yield the matrix factorization.

2.6 Monodromy

It is important to realize that the maps Φ_i and Ψ_i depend on the choice of $i \in \mathbb{Z}$. The effect of changing i is closely related to the concept of monodromy and automorphisms of the triangulated categories involved.

In both $\mathbf{D}(\text{qgr-}A)$ and $\mathbf{D}_{\text{Sg}}^{\text{gr}}(A)$ there is an automorphism of the category generated by

$$M \mapsto M(1). \quad (18)$$

Clearly in terms of coherent sheaves, such a map corresponds to $\mathcal{F} \mapsto \mathcal{F} \otimes \mathcal{O}_X(1)$. This is well-known to correspond to monodromy “around the large radius limit” point, i.e., $B \mapsto B + 1$. In terms of Landau–Ginzburg D-branes, Walcher’s construction shows clearly that this map corresponds to monodromy around the Gepner point.

It is therefore amusing to note that two completely different monodromies are both represented by the same shift in the grading of the A -modules in $\mathbf{D}(\text{gr-}A)$. This is possible, of course, because $\mathbf{D}(\text{qgr-}A)$ and $\mathbf{D}_{\text{Sg}}^{\text{gr}}(A)$ are quite different quotients of $\mathbf{D}(\text{gr-}A)$.

The difference between monodromy around the Gepner point and monodromy around the large radius limit can be determined in terms of monodromy around the “conifold” point. Thus, the above observations should be useful in verifying certain conjectures about monodromy and conifold points as given in [29], for example. We will not pursue this issue here.

Anyway, it is clear from the definitions of Φ_i and Ψ_i that a shift in i simply gives an automorphism of the category induced by monodromy around the large radius limit, or the Gepner point. Such an ambiguity will always be present in maps between the categories of D-branes involved and we are required to make a choice. *From now on, we will choose $i = 0$.* We will see that this choice is consistent with previous statements in the literature.

3 Easy Cases

For the easiest correspondences between matrix factorizations and sheaves it would be nice if we could evade steps 2’ and 3’ in section 2.5. That is, we could consider a free resolution of a module associate to a sheaf and obtain the matrix factorization immediately from its asymptotic form.

To this end, we will prove the following theorem

Theorem 1 *Let $\mathcal{F}$ be the structure sheaf of a projectively normal subvariety of X such that the cohomology groups $H^m(X, \mathcal{F}(r))$ all vanish for $m > 0$ and $r \geq 0$. Let M be the A -module associated with this sheaf. Then if M has no negatively graded part, it lies in $\mathcal{D}_0$. Thus the matrix factorization is obtained directly from the free resolution of M .*

To prove this we follow Orlov [22] again. Let $\pi_0 : \text{gr-}A_{\geq 0} \rightarrow \text{qgr-}A$ be the map given by inclusion into $\text{gr-}A$ followed by the natural projection. Following [30] Orlov shows that this functor has a right adjoint ω_0 which extends to the derived category

$$\mathbf{R}\omega_0 : \mathbf{D}(\text{gr-}A_{\geq 0}) \rightarrow \mathbf{D}(\text{qgr-}A). \quad (19)$$

Furthermore, the image of $\mathbf{R}\omega_0$ is precisely $\mathcal{D}_0$.

Now assume $M \in \text{gr-}A_{\geq 0}$. Then

$$\begin{aligned} M &= \bigoplus_{r=0}^{\infty} M_r \\ &= \bigoplus_{r=0}^{\infty} \text{Hom}_{\text{gr-}A}(A, M(r)). \end{aligned} \tag{20}$$

So

$$\begin{aligned} \omega_0 \pi_0 M &= \bigoplus_{r=0}^{\infty} \text{Hom}_{\text{gr-}A}(A, \omega_0 \pi_0 M(r)) \\ &= \bigoplus_{r=0}^{\infty} \text{Hom}_{\text{qgr-}A}(\pi_0 A, \pi_0 M(r)) \end{aligned} \tag{21}$$

Using derived functors and the definition of sheaf cohomology

$$\begin{aligned} R^j \omega_0 \pi_0 M &= \bigoplus_{r=0}^{\infty} R^j \text{Hom}_{\text{qgr-}A}(\pi_0 A, \pi_0 M(r)) \\ &= \bigoplus_{r=0}^{\infty} H^j(X, \mathcal{F}(r)), \end{aligned} \tag{22}$$

where $\mathcal{F}$ is the sheaf on X associated to M . Assuming $\mathcal{F}$ has no higher cohomology, we see that the complex representing $\mathbf{R}\omega_0 \pi_0 M$ has cohomology only in position zero. From (22) we obtain a homomorphism

$$M \rightarrow \bigoplus_{r=0}^{\infty} H^0(X, \mathcal{F}(r)). \tag{23}$$

Suppose this map is an isomorphism. Then, the map $M \rightarrow \mathbf{R}\omega_0 \pi_0 M$ is a quasi-isomorphism. We would then have proven that M lies in $\mathcal{D}_0$.

So when is the map in (23) an isomorphism? This not true in general but exercise 5.14 in section II of [28] shows that it is true for the structure sheaf of a “projectively normal” variety. This technical condition will be true for most simple examples and can be verified for the examples below. Note, in particular, that a projectively normal variety must be connected.

We may now use theorem 1 to prove some easy equivalences.

3.1 Points

The obvious application is where $\mathcal{F}$ is the skyscraper sheaf $\mathcal{O}_x$ of a point $x \in X$. There is no higher cohomology of a point! The analysis in this case is quite close to section 6.2 of [15]. Let $Y = \text{Proj } B$ be the ambient projective space in which X is embedded. Let I_X be

the homogeneous ideal in B generated by homogeneous functions vanishing on X . Similarly I_x is the homogeneous ideal associated to x . The statement that $x \in X$ corresponds to the inclusion $I_x \supset I_X$.

We may always present a point as a complete intersection in Y , i.e., let $I_x = (f_1, \dots, f_{n-1})$. Clearly $I_X = (W)$. So we have a relation

$$W = \sum_{i=1}^{n-1} f_i g_i, \quad (24)$$

for some $g_i \in B$.

The skyscraper sheaf $\mathcal{O}_x$ corresponds to the A -module $M_x = A/(f_1, \dots, f_{n-1})$. Consider taking a free resolution of this module. By theorem 1 M_x lies in $\mathcal{D}_0$ and, since $\mathcal{O}_x$ is not a trivial sheaf, M_x is not a trivial object in $\mathbf{D}_{\text{Sg}}^{\text{gr}}(A)$.

From what we have just said, this free resolution must be infinite, and the asymptotic form will yield our desired matrix factorization. Following [15] we may write a matrix

$$p = \begin{pmatrix} 0 & p_0 \\ p_1 & 0 \end{pmatrix}, \quad (25)$$

mapping $P_0 \oplus P_1$ on to itself. The matrix factorization condition now becomes $p^2 = W$.

The process of constructing the resolution begins with the functions f_i and analyzes the possible relations between them. Because we have presented the point as a complete intersection or, in other words, because $f_1, \dots, f_{n-1}$ form a “regular sequence”, there will be no “unexpected” relations. We refer to chapter 17 of [31] for more details. Aside from the obvious Koszul relations, we have (24). Because of this, the matrices associated to the maps in the resolution have entries proportional to any of the f_i ’s or g_i ’s. That is

$$p = \sum_{i=1}^{n-1} \pi_i f_i + \bar{\pi}_i g_i, \quad (26)$$

where π_i and $\bar{\pi}_i$ are purely matrices of numbers. The matrix factorization condition then becomes

$$\{\pi_i, \pi_j\} = 0, \quad \{\bar{\pi}_i, \bar{\pi}_j\} = 0, \quad \{\pi_i, \bar{\pi}_j\} = \delta_{ij}. \quad (27)$$

That is, we have the Clifford algebra associated to the Hermitian inner product of $\mathbb{C}^{n-1}$.

For a minimal resolution the matrices π_i and $\bar{\pi}_i$ will have dimension 2^{n-1} and thus our matrices p_0 and p_1 have dimension 2^{n-2} . It is easy to compute these matrices in any example and the 8×8 matrices associated to the quintic threefold were given in [15].

If we consider a hypersurface X in a weighted projective space we need to be careful about the orbifold singularities. If a point lies on a singularity then the matrix factorization will still work just as above but the resulting D-brane will not have the same D-brane charge as a point at a generic smooth point. Instead we will obtain a fractional brane. We will discuss this more in section 5.

It is worth pointing out that another obvious method of producing points is not suitable for the method of theorem 1. Suppose we intersect $n - 2$ generic hyperplanes in $\mathbb{P}^{n-1}$. The intersection of this with $W = 0$ yields n points. Even though these n points are a complete intersection and their higher cohomology is zero, they are not connected and so do not form a projectively normal variety. In this case, (23) is not an isomorphism. Indeed, a free resolution of the associated module in this case is finite and does not produce a matrix factorization.

3.2 Rational Curves

The next easiest case of a sheaf with trivial cohomology is $\mathcal{O}_C$, the structure sheaf of a rational curve $C \cong \mathbb{P}^1$ which lies in X .

If C is a complete intersection in Y , the associated ideal is generated by $n - 2$ equations $f_1, \dots, f_{n-2}$. The fact that this curve lies in X then yields

$$W = \sum_{i=1}^{n-2} f_i g_i, \quad (28)$$

for some polynomials g_i . The construction proceeds in exactly the same way as the previous section. Now the Clifford algebra is associated with $\mathbb{C}^{n-2}$ and our matrix factorization is for matrices of dimension 2^{n-3} .

For example, if X is a 3-fold, the resulting 4×4 matrix factorization can be written

$$p_0 = \begin{pmatrix} f_1 & -g_2 & g_3 & 0 \\ -f_2 & -g_1 & 0 & g_3 \\ f_3 & 0 & -g_1 & g_2 \\ 0 & f_3 & f_2 & f_1 \end{pmatrix}, \quad p_1 = \begin{pmatrix} g_1 & -g_2 & g_3 & 0 \\ -f_2 & -f_1 & 0 & g_3 \\ f_3 & 0 & -f_1 & g_2 \\ 0 & f_3 & f_2 & g_1 \end{pmatrix} \quad (29)$$

The case of the quintic threefold is well-studied. Any line (i.e., curve of degree one) must be the intersection of 3 linear equations and is thus a complete intersection. Thus, the 2875 lines on a generic quintic have 4×4 matrix factorizations. Similarly, any quadric curve lies in a $\mathbb{P}^2 \subset \mathbb{P}^4$ and is thus a complete intersection. So the 609250 quadrics also have this simple matrix factorization.

However, cubic rational curves *cannot* be complete intersections, although they are still projectively normal. If a cubic curve is a complete intersection it would be a plane cubic and thus an elliptic curve. Instead one uses so-called “twisted cubics”. So the cubic rational curves have a slightly more complicated description in terms of matrix factorizations. Having said that, given the presentation of any cubic curve, it is quite easy using a computer package such as Macaulay to compute the corresponding Landau–Ginzburg D-brane. For example, consider the cubic rational curve

$$x_1^2 - x_0 x_2 = x_2^2 - x_1 x_3 = x_1 x_2 - x_0 x_3 = x_4 = 0, \quad (30)$$

in the smooth quintic 3-fold with defining equation

$$x_0^3 x_1 x_2 + x_1^3 x_2^2 + x_1^2 x_2^3 - x_0 x_2^4 - x_0^4 x_3 - x_1^4 x_3 + x_3^4 x_4 + x_4^5. \quad (31)$$

Using Macaulay to compute a resolution in this case gives a 6×6 matrix factorization.

4 Recknagel–Schomerus Branes

Suppose we have an A-series Gepner model. That is, we have a polynomial ring $B = k[x_0, \dots, x_{n-1}]$ where the variables have degrees $d_0, \dots, d_{n-1}$ and a superpotential

$$W = x_0^{l_0} + x_1^{l_1} + \dots + x_{n-1}^{l_{n-1}}, \quad (32)$$

where $d = \sum_i d_i$ and the l_i 's are integral and equal to d/d_i . Define $A = B/(W)$ as above.

View $W = 0$ as an affine variety in k^n . The reason why $\mathbf{D}_{\text{Sg}}^{\text{gr}}(A)$ is not trivial is that this variety has a singularity at the origin. In affine language, the skyscraper sheaf at the origin corresponds to the module $\mathbf{k}$ defined in section 2.4. That is, we have an exact sequence:

$$\bigoplus_{i=0}^{n-1} A(-d_i) \xrightarrow{(x_0 \ x_1 \ \dots \ x_{n-1})} A \longrightarrow \mathbf{k} \longrightarrow 0. \quad (33)$$

We obtain a free resolution by continuing this exact sequence to the left with free modules. This resolution is not finite because of the singularity at the origin.

If we did this resolution in $B = k[x_0, \dots, x_{n-1}]$ we would obtain a finite resolution in terms of the usual Koszul resolution. All the entries in the matrices of this resolution are linear in the x_i 's. When we do the resolution over $A = B/(W)$, we obtain new relations because of the obvious relation

$$\sum_i x_i g_i = W, \quad (34)$$

where $g_i = x_i^{l_i-1}$. We are now in a similar situation to section 3.1. The matrices in the infinite resolution now have entries proportional to x_i or g_i . Then

$$p = \sum_i (x_i \pi_i + g_i \bar{\pi}_i), \quad (35)$$

where π_i and $\bar{\pi}_i$ are matrices of numbers. The matrix factorization condition then produces the Clifford algebra (27) associated to $\mathbb{C}^n$. Thus p_0 and p_1 are matrices of dimension 2^{n-1} . This is exactly the construction used in [15] to produce the tensor product D-branes of the Gepner model. That is, we claim that $\mathbf{k}$ represents the tensor product D-brane

$$M_1(x_0) \otimes M_1(x_1) \otimes \dots \otimes M_1(x_{n-1}), \quad (36)$$

where $M_a(x)$ represents the matrix factorization of the minimal model $x^a \cdot x^{m-a} = x^m$.

It was demonstrated in [15] that these tensor product D-branes correspond to the RS D-branes of [9] constructed directly from the Gepner model. An RS D-brane is denoted by⁶ $(L_0, L_1, \dots, L_{n-1})_r$. The result of [15] is that the D-brane $(L_0, L_1, \dots, L_{n-1})_0$ is equivalent to the matrix factorization

$$M_{L_0+1}(x_0) \otimes M_{L_1+1}(x_1) \otimes \dots \otimes M_{L_{n-1}+1}(x_{n-1}). \quad (37)$$

⁶Our subscript will be r rather than the more conventional M to be consistent with the notation in the rest of this paper.

Thus we have shown the correspondence of D-branes

$$(0, 0, \dots, 0)_0 \cong \mathbf{k}. \quad (38)$$

Replacing x_j by $x_j^{L_j+1}$ in the free resolution above, it is easy to see that we have a more general correspondence

$$(L_0, L_1, \dots, L_{n-1})_0 \cong \frac{A}{(x_0^{L_0+1}, x_1^{L_1+1}, \dots, x_{n-1}^{L_{n-1}+1})}. \quad (39)$$

From Walcher's construction [25], it is easy to see that shifting r in the RS D-branes is equivalent to shifting the grading of the module. Thus we have the following correspondence for an arbitrary RS D-brane:

$$(L_0, L_1, \dots, L_{n-1})_r \cong \frac{A}{(x_0^{L_0+1}, x_1^{L_1+1}, \dots, x_{n-1}^{L_{n-1}+1})}(r). \quad (40)$$

This completes step 1 of section 2.5. We have found an A -module whose free resolution corresponds to the desired matrix factorization associated to any Recknagel–Schomerus D-brane.

4.1 Type $(0, 0, \dots, 0)_r$

In order to proceed let us focus on some specific D-branes. First consider $(0, 0, \dots, 0)_0 \cong \mathbf{k}$. Clearly $\mathbf{k}$ has only a degree 0 part with respect to the grading and so step 2 of section 2.5 is trivial — $\mathbf{k}$ is already in $\mathbf{D}(\text{gr-}A_{\geq 0})$.

Suppose, for a moment, that step 3 were trivial too, i.e., that $\mathbf{k}$ lay in $\mathcal{T}_0$. This would mean that our resulting sheaf in the large radius interpretation would be the sheaf on X corresponding to the A -module $\mathbf{k}$. But $\mathbf{k}$ is a torsion sheaf and so, by Serre's construction, corresponds to the zero sheaf. We know that $(0, 0, \dots, 0)_0$ is a nontrivial D-brane will results in a contradiction.

To perform step 3 we need a distinguished triangle

$$\begin{array}{ccc} p & \xrightarrow{[1]} & t \\ & \searrow & \swarrow \\ & \mathbf{k} & \end{array} \quad (41)$$

where p is in $\mathcal{P}_{\geq 0}$ and t is in $\mathcal{T}_0$. From the definition of the semiorthogonal decomposition, and the definition of $\mathcal{P}_{\geq 0}$, t must satisfy⁷

$$\text{Ext}^m(t, A(r)) = 0, \quad \text{for all } m \text{ and for all } r \leq 0. \quad (42)$$

⁷From now on, any Hom or Ext appearing without a subscript will refer to the (derived) category of graded A -modules.

From the long exact sequence of Ext's associated to (41) we obtain

$$\mathrm{Ext}^m(p, A(r)) = \mathrm{Ext}^m(\mathbf{k}, A(r)), \quad \text{for all } m \text{ and for all } r \leq 0. \quad (43)$$

Now, A is an “AS-Gorenstein” ring since

$$\mathbf{R} \mathrm{Hom}_A(\mathbf{k}, A) = \mathbf{k}[-n+1]. \quad (44)$$

This follows from Serre duality and the Calabi–Yau condition on X . We refer to lemma 2.11 in [22] for a proof of this statement. It follows that

$$\mathrm{Ext}^{n-1}(\mathbf{k}, A) = k, \quad (45)$$

with all the other Ext groups vanishing in (43). Thus we may satisfy (43), and therefore (42), by setting $p = A[n-1]$.

The distinguished triangle (41) can also be used to perform step 4. We need to project the object t into in $\mathbf{D}(\mathrm{qgr}\text{-}A) = \mathbf{D}(X)$. But $\mathbf{D}(\mathrm{qgr}\text{-}A)$ is obtained by quotienting by torsion modules and $\mathbf{k}$ is itself a torsion module. Thus, using the definition of a quotient triangulated category (and shuffling around the $[1]$ in (41)), we see that t is equivalent to $p[-1] = A[n-2]$ in $\mathbf{D}(\mathrm{qgr}\text{-}A)$. It is basic to the construction of sheaves that A as an A -module corresponds to the structure sheaf $\mathcal{O}_X$. We then obtain the correspondence

$$(0, 0, \dots, 0)_0 \cong \mathcal{O}_X[n-2]. \quad (46)$$

In the case of the quintic threefold, this correspondence has been known (up to the shift) for some time [10]. Now we see this result is completely general.

Next let us consider $(0, 0, \dots, 0)_{-1} \cong \mathbf{k}(-1)$. This is very similar to the above case of $(0, 0, \dots, 0)_0$. Again the projection to $\mathbf{D}(\mathrm{gr}\text{-}A_{\geq 0})$ is trivial. Now, we require p to satisfy $\mathrm{Ext}^{n-1}(p, A(-1)) = k$. This could be satisfied by setting $p = A(-1)[n-1]$. However, we also require $\mathrm{Ext}^m(p, A) = 0$ for all m and this choice would break this latter condition.

Recall that $\mathrm{Hom}_A(A, A) = A$ as an A -module. Thus, in the category of graded A -modules, $\mathrm{Hom}(A, A(r)) = A_r$ (i.e., the vector space of elements of degree r in the algebra A) with all higher Ext's vanishing. We claim one should use the following complex for p :

$$A(-1) \xrightarrow[\boxed{1-n}]{\begin{pmatrix} x_0 \\ x_1 \\ \vdots \\ x_{\alpha_1-1} \end{pmatrix}} A^{\oplus \alpha_1}, \quad (47)$$

where $x_0, x_1, \dots, x_{\alpha_1-1}$ are the variables of degree one in the ring B . The $1-n$ in a box denotes the position in the complex. Applying $\mathrm{Hom}(-, A(-1))$ we see that $\mathrm{Ext}^{n-1}(p, A(-1)) = k$. Applying $\mathrm{Hom}(-, A)$ to this complex, the two terms in the complex cancel and we obtain $\mathrm{Ext}^{n-1}(p, A) = 0$ as desired. It is easy to show that all other Ext's vanish as necessary justifying our claim.

Thus, in general, the D-brane $(0, 0, \dots, 0)_{-1}$ corresponds to the following complex of sheaves

$$\mathcal{O}_X(-1) \xrightarrow[\boxed{2-n}]{\begin{pmatrix} x_0 \\ x_1 \\ \vdots \\ x_{\alpha_1-1} \end{pmatrix}} \mathcal{O}_X^{\oplus \alpha_1}. \quad (48)$$

Suppose all the variables in B are of degree 1, i.e., we have a non-weighted projective space. Then d , the degree of the superpotential, equals n . On the corresponding $\mathbb{P}^{n-1}$ we have the Euler exact sequence

$$0 \longrightarrow \mathcal{O} \xrightarrow{\begin{pmatrix} x_0 \\ x_1 \\ \vdots \\ x_{n-1} \end{pmatrix}} \mathcal{O}(1)^n \longrightarrow \mathcal{T} \longrightarrow 0, \quad (49)$$

where $\mathcal{T}$ is the tangent sheaf of $\mathbb{P}^{n-1}$. It follows that $(0, 0, \dots, 0)_{-1}$ corresponds to $\mathcal{T}(-1)[n-3]$ restricted to X .

Let Ω^k denote the k th exterior power of the cotangent bundle of $\mathbb{P}^{n-1}$ restricted to X . Since $\Omega^{n-1} \cong \mathcal{O}_X(-n)$, it follows that $\mathcal{T}(-1)[n-3] = \Omega^{n-2}(n-1)[n-3]$.

In order to see the general picture (for arbitrary weights) for $(0, \dots, 0)_r = \mathbf{k}(r)$ for any r , it is instructive to consider the cases where $r > 0$. Consider constructing a free resolution $C \rightarrow \mathbf{k}$, where C is an infinite complex of free A -modules. This leads to a distinguished triangle in $\mathbf{D}(\text{gr-}A)$:

$$\begin{array}{ccc} C(r)_{\geq 0} & \xrightarrow{[1]} & C(r)_{< 0} \\ & \nwarrow \quad \nearrow & \\ & \mathbf{k}(r) & \end{array} \quad (50)$$

where $C(r)_{\geq 0}$ is obtained from $C(r)$ by deleting from the complex any $A(a)$ for $a > 0$. Similarly $C(r)_{< 0}$ is obtained from $C(r)$ by deleting all the $A(a)$'s for $a \leq 0$. Clearly $C(r)_{< 0}$ is a finite length complex.

$C(r)_{\geq 0}$ is trivially in $\mathbf{D}(\text{gr-}A_{\geq 0})$. Less obviously, it is also in $\mathcal{T}_0$ as can be seen as follows. Since $\text{Ext}^m(C(r)_{< 0}, A(s)) = 0$ for any m and any $s \leq 0$, it follows from (50) that $\text{Ext}^m(C(r)_{\geq 0}, A(s)) = \text{Ext}^m(\mathbf{k}(r), A(s))$ for any m and any $s \leq 0$. But, given that $r > 0$, we see from (44) that the latter is zero. It follows that $C(r)_{\geq 0}$ is in $\mathbf{D}(\text{gr-}A_{\geq 0})$. Finally we may use (50) again to see that $C(r)_{\geq 0}$ is equivalent to $C(r)_{< 0}[1]$ in $\mathbf{D}(\text{qgr-}A)$.

We have therefore shown that the matrix factorization corresponding to $\mathbf{k}(r)$ is equivalent to the finite complex $C(r)_{< 0}[1]$ in $\mathbf{D}(\text{qgr-}A)$.

To simply matters further, suppose $0 < r < d$. In this case, the complex $C(r)_{< 0}$ obviously never contains any part of the free resolution that gets to “see” the effects of the condition $W = 0$. This means that we may as well resolve $\mathbf{k}(r)$ as a B -module, rather than an A -module. This is a straight-forward finite Koszul resolution.

r	$(0, 0, 0, 0, 0)_r$
0	$\mathcal{O}_X \longrightarrow 0 \longrightarrow 0$
1	$\mathcal{O}_X(1)$
2	$\mathcal{O}_X(1)^{\oplus 2} \xrightarrow{(x_3 \ x_4)} \mathcal{O}_X(2)$
3	$\mathcal{O}_X(1) \longrightarrow \mathcal{O}_X(2)^{\oplus 2} \oplus \mathcal{O}_X(1)^{\oplus 3} \longrightarrow \mathcal{O}_X(3)$
4	$\mathcal{O}_X(2) \oplus \mathcal{O}_X(1)^{\oplus 6} \longrightarrow \mathcal{O}_X(3)^{\oplus 2} \oplus \mathcal{O}_X(2)^{\oplus 3} \longrightarrow \mathcal{O}_X(4)$
5	$\mathcal{O}_X(1)^{\oplus 3} \longrightarrow \mathcal{O}_X(3) \oplus \mathcal{O}_X(2)^{\oplus 6} \oplus \mathcal{O}_X(1)^{\oplus 3} \longrightarrow \mathcal{O}_X(4)^{\oplus 2} \oplus \mathcal{O}_X(3)^{\oplus 3} \longrightarrow \mathcal{O}_X(5)$
6	$\mathcal{O}_X(2)^{\oplus 3} \oplus \mathcal{O}_X(1)^{\oplus 6} \longrightarrow \mathcal{O}_X(4) \oplus \mathcal{O}_X(3)^{\oplus 6} \oplus \mathcal{O}_X(2)^{\oplus 3} \longrightarrow \mathcal{O}_X(5)^{\oplus 2} \oplus \mathcal{O}_X(4)^{\oplus 3} \longrightarrow \mathcal{O}_X(6)$
7	$\mathcal{O}_X(1)^{\oplus 3} \longrightarrow \mathcal{O}_X(3)^{\oplus 3} \oplus \mathcal{O}_X(2)^{\oplus 6} \oplus \mathcal{O}_X(1) \longrightarrow \mathcal{O}_X(5) \oplus \mathcal{O}_X(4)^{\oplus 6} \oplus \mathcal{O}_X(3)^{\oplus 3} \longrightarrow \mathcal{O}_X(6)^{\oplus 2} \oplus \mathcal{O}_X(5)^{\oplus 3} \longrightarrow \mathcal{O}_X(7)$

Table 1: The case $\mathbb{P}^4_{\{2,2,2,1,1\}}$. The final term in the complex is always in position -1 .

This makes computing the geometrical interpretation of the D-brane $\mathbf{k}(r)$ very easy. In the case of a non-weighted projective space (so $d = n$) we obtain

$$(0, 0, \dots, 0)_r \cong \Omega^{r-1}(r)[r], \quad (51)$$

for $0 < r < n$. For the quintic threefold, putting $n = d = 5$, we recover the well-known sequence of sheaves as discussed in [10], for example. Given (46) and (9) we know the geometrical interpretation for $(0, 0, \dots, 0)_r$ for all r .

The case of weighted projective space is more complicated although it is always true that

$$(0, 0, \dots, 0)_1 \cong \mathcal{O}_X(1)[1]. \quad (52)$$

Let us consider, as an example, the familiar case of weighted $\mathbb{P}^4$ with weights $\{2, 2, 2, 1, 1\}$ and $d = 8$. It is easy to compute the complexes of sheaves corresponding to $(0, 0, \dots, 0)_r$ and we list the result in table 1.

Note that the complex corresponding to $(0, 0, \dots, 0)_2$ has nontrivial cohomology in both positions. Thus this complex cannot possibly be equivalent to a single sheaf. This is in contrast with the result (51) for the non-weighted case. At position -1 , the cohomology is equal to $\mathcal{O}_S(2)[1]$, where $\mathcal{O}_S$ is the structure sheaf of the $\mathbb{Z}_2$ -orbifold singularity $x_3 = x_4 = 0$.

The sequence of complexes in table 1 shows how $\mathcal{O}_X$ behaves under monodromy around the Landau–Ginzburg point in the moduli space as discussed in [32], for example. The same result, using Fourier–Mukai transforms, has also been observed recently in [33].

4.2 General Torsion Modules

It is very straight-forward to build the general RS-brane $(L_0, L_1, \dots)_r$ from the branes $\mathbf{k}(r) \cong (0, 0, \dots)_r$ of the preceding section. Consider the following short exact sequence:

$$0 \longrightarrow \frac{k[x]}{(x)}(-1) \xrightarrow{x} \frac{k[x]}{(x^2)} \longrightarrow \frac{k[x]}{(x)} \longrightarrow 0. \quad (53)$$

This shows that $k[x]/(x^2)$ is an extension coming from $\text{Ext}^1(k[x]/(x), (k[x]/(x))(-1))$. Using the concept of this extension, it is straight-forward to build any RS D-brane as a finite sequence of extensions of a collection of branes of the type $(0, \dots, 0)_r$ considered in the previous section.

For example, we immediately have a distinguished triangle

$$\begin{array}{ccc} (L_0, L_1, \dots, L_{n-1})_0 & \xrightarrow{[1]} & (0, L_1, \dots, L_{n-1})_{-L_0} \\ & \nwarrow & \nearrow x_0^{L_0+1} \\ & (L_0 + 1, L_1, \dots, L_{n-1})_0 \end{array} \quad (54)$$

This observation leads to a systematic picture of the relations between various RS D-branes.

It is quite easy in this picture to confirm the finiteness of the number of RS D-branes. Suppose we have a D-brane of the form (40) where one of the $(L_i + 1)$'s is equal to l_i , the degree of the corresponding variable in the superpotential W . Then we may use the superpotential relation to remove this monomial from the set of generators of the ideal in the denominator of (40). Thus, if we perform a free resolution of this module, the variable x_i never appears in the maps. Thus, this free resolution can equally be performed over the ring B and is therefore finite. It follows that the module is zero in $\mathbf{D}_{\text{sg}}^{\text{gr}}(A)$. This also leads to the brane-anti-brane relation

$$(L_0, L_1, \dots, L_i, \dots, L_{n-1})_0 \cong (L_0, L_1, \dots, L_{l_i-L_i-2}, \dots, L_{n-1})_{-L_{i-1}}[1], \quad (55)$$

and so we need only consider the range $0 \leq L_i \leq (l_i - 2)/2$ (as is also clear from the original construction [9]).

Another relation we obtain from triangles of the form (54) is

$$(1, 0, \dots, 0)_1 = \text{Cone}((0, 0, \dots, 0)_1[-1] \rightarrow (0, 0, \dots, 0)_0). \quad (56)$$

Using the results of section 4.1, this yields an equivalence

$$(1, 0, \dots, 0)_1 \cong \text{Cone}(O_X(1) \rightarrow O_X[n-2]). \quad (57)$$

For the quintic 3-fold, with $n = 5$, this result was observed in [3] and the stability of this “exotic” state was discussed in [6]. Now we see that the relationship (57) is true in general for the whole class of models we are studying in this paper.

Actually it is natural to consider the set of D-branes that can be finitely generated from the set $\mathbf{k}(r) \cong (0, 0, \dots)_r$. This is precisely the set of modules which are *torsion* modules, i.e., modules which are finite-dimensional as vector spaces. This set includes all the RS-branes but includes many more. Let $\mathcal{S}$ denote the full subcategory of $\mathbf{D}(\text{gr-}A)$ corresponding to bounded complexes of torsion modules. For example, the module

$$M = \frac{A}{(x_0^2, x_1^2, x_0x_1, x_2, x_3, x_4)} \quad (58)$$

is in $\mathcal{S}$ but is not an RS D-brane. However, it can be built from RS D-branes:

$$M = \text{Cone}((0, 1, 0, 0, 0)[-1] \rightarrow (0, 0, 0, 0, 0)_{-1}). \quad (59)$$

The set $\mathcal{S}$ is of interest for the following reason. For simplicity let us focus on the case of non-weighted projective spaces. In that case, it is known from the work of Beilinson [34] that the set of sheaves in (51) form a complete exceptional collection on the ambient $\mathbb{P}^{n-1}$. That is, they generate the whole derived category of coherent sheaves on $\mathbb{P}^{n-1}$. Since $\mathcal{S}$ is also precisely the category generated by these objects, we observe that *the image of $\mathcal{S}$ in $\mathbf{D}(\text{qgr-}A)$ is precisely the subset of D-branes on X that arise as complexes of restrictions of sheaves on the ambient $\mathbb{P}^{n-1}$.*⁸

5 Quotient Singularities

In general, a weighted projective space has quotient singularities. The hypersurface $W = 0$ may also therefore have quotient singularities. Thus far we have been a little careless about our language of sheaves on X when we have such singularities.

More properly we should define the quotient *stack* $\text{Proj } A$ as $[(\text{Spec } A \setminus 0)/k^*]$ on which it is easier to define coherent sheaves.

The problem is seen by a simple example copied from [35]. Let us consider sheaves on the weighted projective space $\mathbb{P}_{\{1,1,2\}}^2$ and let B be its homogeneous coordinate ring $k[x_0, x_1, x_2]$, where x_2 has degree 2. This space has a $\mathbb{Z}_2$ -quotient singularity at $[0, 0, 1]$. Consider first the module

$$M = \frac{B}{(x_0, x_1)}. \quad (60)$$

The stalk of the associated sheaf $\widetilde{M}$ at the point $[x_0, x_1, x_2] = [a_0, a_1, a_2]$ is obtained by localizing this module on the homogeneous ideal of functions vanishing at this point and restricting to elements of degree zero. This localization is zero at every point except $[0, 0, 1]$. The stalk at $[0, 0, 1]$ is k and thus this is the skyscraper sheaf of $[0, 0, 1]$.

Now consider

$$M(1) = \frac{B}{(x_0, x_1)}(1). \quad (61)$$

⁸The sheaves in the complex are restrictions of sheaves on $\mathbb{P}^{n-1}$ but the morphisms in the complex need not lift to morphisms on $\mathbb{P}^{n-1}$. Thus it would not be correct to assert that $\mathcal{S}$ is the image of $\mathbf{D}(\mathbb{P}^{n-1})$ in $\mathbf{D}(X)$.

This is the same as M except that when we localize at $[0, 0, 1]$ we get zero due to the choice of gradings of the variables. That is, $\widetilde{M}(1)$ is the zero sheaf. So Serre's correspondence between graded modules modulo torsion and coherent sheaves is not strictly correct for weighted projective spaces.

Instead of coherent sheaves on $X = \text{Proj } A$, one should consider coherent sheaves on $\text{Spec } A$ which are equivariant under the k^* action producing the weighted projective space. In [22] it was shown that the category of such objects is equivalent to $\text{qgr-}A$ as required for our program.

Suppose a weighted projective space has an orbifold singularity locally of the form k^{n-2}/G for some discrete group $G \subset \text{SL}(n-2, \mathbb{Z})$. Then a k^* -equivariant sheaf on $\text{Spec } A$ is locally modeled by a G -equivariant sheaf on k^{n-2} .

In the case of $\mathbb{P}_{\{1,1,2\}}^2$ analyzed above, it is clear how to interpret our sheaves now. Consider the skyscraper sheaf of the origin in k^2 . A $\mathbb{Z}_2$ action on k^2 which fixes the origin may act as ± 1 on such a sheaf. The resulting quotient sheaf will be associated to the modules M or $M(1)$ respectively. For a longer discussion of this and how it fits into the language of stacks we refer, for example, to [36].

This is exactly the setting for the *McKay correspondence*. Locally the McKay correspondence asserts that the derived category of G -equivariant sheaves on k^{n-2}/G is equivalent to the derived category of coherent sheaves on a crepant resolution of this singularity, assuming such a resolution exists. A necessary condition for the existence of such a resolution is that $G \subset \text{SL}(n-2, \mathbb{Z})$. Mathematically the McKay correspondence has only been proven in certain cases (see [37], and [38] for a review). String theory essentially “proves” the McKay correspondence in all cases since blowing up the orbifold is a change in $B + iJ$ which cannot effect the B-model. Thus, B-type D-branes are unaffected by the process. See [39] for further details.

Let $\widetilde{X}$ be a smooth crepant resolution of X , assuming one exists. A global form of the McKay correspondence, which we will assume here, states an equivalence between the derived category of sheaves on X (viewed as a stack) and the derived category of sheaves on $\widetilde{X}$. So, we arrive at the statement that *the category $D\text{Gr}B(W)$ of matrix factorizations is equivalent to the category $\mathbf{D}(\widetilde{X})$* .

Actually the category $\mathbf{D}(\widetilde{X})$ may be constructed directly following the construction of Cox [35]. $\widetilde{X}$ is now a hypersurface in some toric variety T obtained by blowing up the weighted projective space. A is now a multigraded algebra derived from the homogeneous coordinates of T . We construct some ideal⁹ J_Σ derived from the toric fan Σ associated to T . Then

$$\mathbf{D}(\widetilde{X}) \cong \frac{\mathbf{D}(\text{gr-}A)}{\mathcal{J}_\Sigma}, \quad (62)$$

where $\mathcal{J}_\Sigma$ is the triangulated category generated by J_Σ and all its multigraded twists. This construction seems to fit very naturally into Orlov's picture of equivalences between categories but we will not pursue it further here.

⁹Denoted B by Cox.

5.1 $\mathbb{P}_{\{2,2,2,1,1\}}^4$

Let us consider how all this works in the example $\mathbb{P}_{\{2,2,2,1,1\}}^4$ studied in [40]. Some aspects of D-branes and matrix factorizations for this case have already been analyzed in [20]. This model has a $\mathbb{Z}_2$ singularity along the curve C given by $z_3 = z_4 = 0$. C is a curve of genus 3. Let D be the exceptional divisor $C \times \mathbb{P}^1$ one obtains when this curve is blown up.

Let us remind ourselves of the McKay correspondence for the simplest case $k^2/\mathbb{Z}_2$. Start with the skyscraper sheaf of the origin in k^2 . We may divide this by $\mathbb{Z}_2$ equivariantly in two ways according to whether the $\mathbb{Z}_2$ acts as $+1$ or -1 on the fibre. We denote these two $\mathbb{Z}_2$ -equivariant sheaves $\mathcal{O}_0^+$ and $\mathcal{O}_0^-$. The resolution of $k^2/\mathbb{Z}_2$ has $E = \mathbb{P}^1$ as the exceptional divisor. Let $\mathcal{O}_E$ denote the structure sheaf of E extended by zero over the total space of the resolution. The McKay correspondence then maps

$$\begin{aligned}\mathcal{O}_0^+ &\mapsto \mathcal{O}_E \\ \mathcal{O}_0^- &\mapsto \mathcal{O}_E(-1)[1].\end{aligned}\tag{63}$$

Extending this to the 3-fold in question, the sheaf $\mathcal{O}_D$, corresponding to a 4-brane wrapping D , will correspond to the $\mathbb{Z}_2$ -invariant structure sheaf of C in $\mathbb{P}_{\{2,2,2,1,1\}}^4$. That is, we consider the module $M = A/(x_3, x_4)$.

We may now follow the procedure of section 2.5 to yield the corresponding matrix factorization. We have already done step 1', and step 2' is trivial since M is already in $\mathbf{D}(\text{gr-}A_{\geq 0})$. The nontrivial part of the analysis comes in step 3'. A computation with Macaulay yields

$$\begin{aligned}\text{Ext}^2(\mathbf{k}(-2), M) &= k \\ \text{Ext}^3(\mathbf{k}(-1), M) &= k^2 \\ \text{Ext}^4(\mathbf{k}, M) &= k,\end{aligned}\tag{64}$$

showing that M does not lie in $\mathcal{D}_0$. Instead one builds the following complex:

$$M'' = \left(\mathbf{k}[-3]^{\oplus 3} \xrightarrow{(x_0 \ x_1 \ x_2)} \mathbf{k}(-2)[-2] \xrightarrow[f]{\quad} M \right), \tag{65}$$

where f generates $\text{Ext}^2(\mathbf{k}(-2), M)$. M'' satisfies $\text{Ext}^m(\mathbf{k}(-r), M'')$ for all m and all $r \geq 0$ and so lies in $\mathcal{D}_0$. It is easy to show that M has a finite free resolution and so does not contribute to the matrix factorization. This results in the correspondence:

$$\mathcal{O}_D \cong \text{Cone} \left(\mathbf{k}[-4]^{\oplus 3} \xrightarrow{(x_0 \ x_1 \ x_2)} \mathbf{k}(-2)[-3] \right).$$

(66)

In section 4 we identified the D-branes $\mathbf{k}(r) = (0, 0, \dots, 0)_r$ as specific matrix factorizations. In our case, with $n = 5$, they are given by 16×16 matrices. These may be substituted into the above equation to yield an explicit 64×64 matrix factorization associated to $\mathcal{O}_D$.

Now consider $M(1)$. The $\mathbb{Z}_2 \subset k^*$ that acts as -1 on x_3 and x_4 , will act as -1 on this module. So this is the $\mathbb{Z}_2$ -anti-invariant sheaf supported on C . It follows from (63) and (66) that

$$\mathcal{O}_D(0, -1) \cong \text{Cone}\left(\mathbf{k}(1)[-5]^{\oplus 3} \xrightarrow{(x_0 \ x_1 \ x_2)} \mathbf{k}(-1)[-4]\right). \quad (67)$$

yielding a 64×64 matrix factorization for this sheaf.

Note that we have a bigrading for the sheaf on the left of (67). This is because $h^{1,1} = 2$ for the resolved manifold and so we have two directions in the Kähler moduli space. If we denote a twisted sheaf by $\mathcal{F}(a, b)$, then monodromy around the large radius limit of the weighted projective space itself will increase a while monodromy around the large radius limit of the exceptional set will increase b . See [32] for more on these monodromies.

As a further example, suppose we want to find the structure sheaf of the subspace $C \times \{p\}$ of the exceptional set $C \times \mathbb{P}^1$. Let p have homogeneous coordinates $[y_0, y_1]$. One can then show that

$$\mathcal{O}_{C \times \{p\}} = \text{Cone}\left(\mathcal{O}_D(0, -1) \xrightarrow{y_1 x_3 - y_0 x_4} \mathcal{O}_D\right). \quad (68)$$

Therefore this sheaf may be built by coning the above matrices to form a factorization using 128×128 matrices.

6 Discussion

We have demonstrated a completely systematic way of translating between the geometric language of vector bundles or sheaves and the language of matrix factorizations. At first sight one might consider it an advantage to try to compute with matrix factorizations. After all, the concept of a matrix factorization would appear to be more straight-forward than the derived category of coherent sheaves. However, we have discovered that the matrices for geometrical objects rapidly become large — we observed a 128×128 matrix factorization for a 2-brane in section 5.1. It is also very awkward to compute the space of morphisms between two matrix factorizations. So we believe that there is no computational advantage in using matrix factorizations over coherent sheaves.

It is interesting to observe that, for 3-folds, the complete intersection rational curves appear to correspond to the lowest-dimensional matrices. In the case of the quintic threefold these 4×4 matrices corresponds to the 2875 lines and 609250 quadrics. One can easily show that 2×2 matrix factorizations are impossible for a smooth 3-fold and we have not observed any 3×3 matrix factorizations.

The construction of the category of B-type D-branes in the paper has some very striking features which appear to make it a very natural setting for the phase picture of Calabi–Yau manifolds. There is a single category — namely the derived category of graded A -modules that seems to serve as the “master” category over the whole moduli space. The

Landau–Ginzburg phase and the large radius phase have D-branes described by particular quotient categories of this master category. Monodromy around the limit point in each phase corresponds to shifting the grade by one in the master category.

The semiorthogonal decompositions give a very explicit method of taking the various quotients of the master category and allows one to translate between the different phases with surprising ease.

Obtaining the general phase picture in toric geometry would seem to simply be a matter of extending Orlov’s results to multiply-graded algebras. We will pursue this elsewhere.

Another obvious extension of this work involves analyzing stability for D-branes in Landau–Ginzburg theories. Although there has been some work on this subject [25], there remains much to be discovered. There is a general belief that stability conditions should simplify at the limit of point of each phase. Therefore there might be a relationship between stability and Orlov’s picture of semiorthogonal decompositions.

Acknowledgments

I wish to thank T. Bridgeland, D. Orlov, A. Roy and J. Walcher for useful discussions. The author is supported by NSF grants DMS–0301476 and DMS–0606578.

References

- [1] E. Witten, *Phases of $N = 2$ Theories in Two Dimensions*, Nucl. Phys. **B403** (1993) 159–222, hep-th/9301042.
- [2] P. S. Aspinwall, B. R. Greene, and D. R. Morrison, *Calabi–Yau Moduli Space, Mirror Manifolds and Spacetime Topology Change in String Theory*, Nucl. Phys. **B416** (1994) 414–480.
- [3] M. R. Douglas, *D-Branes, Categories and $N=1$ Supersymmetry*, J. Math. Phys. **42** (2001) 2818–2843, hep-th/0011017.
- [4] M. R. Douglas, *Dirichlet Branes, Homological Mirror Symmetry, and Stability*, in “Proceedings of the International Congress of Mathematicians: Beijing 2002”, vol. III, pages 395–408, Higher Education Press, Beijing, 2003, math.AG/0207021.
- [5] P. S. Aspinwall and A. E. Lawrence, *Derived Categories and Zero-Brane Stability*, JHEP **08** (2001) 004, hep-th/0104147.
- [6] P. S. Aspinwall and M. R. Douglas, *D-Brane Stability and Monodromy*, JHEP **05** (2002) 031, hep-th/0110071.
- [7] D. Gepner, *Exactly Solvable String Compactifications on Manifolds of $SU(N)$ Holonomy*, Phys. Lett. **199B** (1987) 380–388.

- [8] C. Vafa and N. Warner, *Catastrophes and the Classification of Conformal Theories*, Phys. Lett **218B** (1989) 51–58.
- [9] A. Recknagel and V. Schomerus, *D-branes in Gepner Models*, Nucl. Phys. **B531** (1998) 185–225, hep-th/9712186.
- [10] I. Brunner, M. R. Douglas, A. Lawrence, and C. Römelsberger, *D-branes on the Quintic*, JHEP **08** (2000) 015, hep-th/9906200.
- [11] A. Kapustin and Y. Li, *D-branes in Landau-Ginzburg Models and Algebraic Geometry*, JHEP **12** (2003) 005, hep-th/0210296.
- [12] A. Kapustin and Y. Li, *D-branes in Topological Minimal models: The Landau–Ginzburg Approach*, JHEP **07** (2004) 045, hep-th/0306001.
- [13] I. Brunner, M. Herbst, W. Lerche, and B. Scheuner, *Landau-Ginzburg realization of open string TFT*, hep-th/0305133.
- [14] C. I. Lazaroiu, *On the Boundary Coupling of Topological Landau–Ginzburg Models*, JHEP **05** (2005) 037, hep-th/0312286.
- [15] S. K. Ashok, E. Dell’Aquila, and D.-E. Diaconescu, *Fractional Branes in Landau–Ginzburg Orbifolds*, Adv. Theor. Math. Phys. **8** (2004) 461–513, hep-th/0401135.
- [16] I. Brunner, M. Herbst, W. Lerche, and J. Walcher, *Matrix Factorizations and Mirror Symmetry: The Cubic Curve*, hep-th/0408243.
- [17] M. R. Douglas, S. Govindarajan, T. Jayaraman, and A. Tomasiello, *D-branes on Calabi–Yau Manifolds and Superpotentials*, Commun. Math. Phys. **248** (2004) 85–118, hep-th/0203173.
- [18] K. Hori and J. Walcher, *F-term Equations Near Gepner Points*, JHEP **01** (2005) 008, hep-th/0404196.
- [19] K. Hori and J. Walcher, *D-branes from Matrix Factorizations*, Comptes Rendus Physique **5** (2004) 1061–1070, hep-th/0409204.
- [20] I. Brunner and M. R. Gaberdiel, *Matrix Factorisations and Permutation Branes*, JHEP **07** (2005) 012, hep-th/0503207.
- [21] H. Enger, A. Recknagel, and D. Roggenkamp, *Permutation Branes and Linear Matrix Factorisations*, JHEP **01** (2006) 087, hep-th/0508053.
- [22] D. Orlov, *Derived Categories of Coherent Sheaves and Triangulated Categories of Singularities*, math.AG/0503632.
- [23] D. Orlov, *Triangulated Categories of Singularities and D-Branes in Landau–Ginzburg Orbifolds*, Proc. Steklov Inst. Math. **246** (2004) 227–248, math.AG/0302304.

- [24] A. Kapustin and Y. Li, *Topological Correlators in Landau–Ginzburg Models with Boundaries*, Adv. Theor. Math. Phys. **7** (2004) 727–749, hep-th/0305136.
- [25] J. Walcher, *Stability of Landau–Ginzburg Branes*, J. Math. Phys. **46** (2005) 082305, hep-th/0412274.
- [26] D. Eisenbud, *Homological Algebra on a Complete Intersection, with an Application to Group Representations*, Trans. Amer. Math. Soc. **260** (1980) 35–64.
- [27] J. P. Serre, *Modules projectifs et espace fibrés à fibre vectorielle*, Séminaire Dubreil–Pisot **23** (1958) 531–543.
- [28] R. Hartshorne, *Algebraic Geometry*, Graduate Texts in Mathematics **52**, Springer-Verlag, 1977.
- [29] P. S. Aspinwall, R. L. Karp, and R. P. Horja, *Massless D-branes on Calabi-Yau threefolds and monodromy*, Commun. Math. Phys. **259** (2005) 45–69, hep-th/0209161.
- [30] M. Artin and D. Mumford, *Some Elementary Results of Unirational Varieties which are not Rational*, Proc. London Math. Soc. **25** (1972) 75–95.
- [31] D. Eisenbud, *Commutative Algebra With a View Towards Algebraic Geometry*, Graduate Texts in Mathematics **150**, Springer, 2004.
- [32] P. S. Aspinwall, *Some Navigation Rules for D-brane Monodromy*, J. Math. Phys. **42** (2001) 5534–5552, hep-th/0102198.
- [33] A. Canonaco and R. L. Karp, *Derived Autoequivalences and a Weighted Beilinson Resolution*, to appear.
- [34] A. A. Beilinson, *Coherent Sheaves on $\mathbb{P}^n$ and Problems in Linear Algebra*, Func. Anal. Appl. **12** (1978) 214–216.
- [35] D. A. Cox, *The Homogeneous Coordinate Ring of a Toric Variety*, J. Algebraic Geom. **4** (1995) 17–50, alg-geom/9210008.
- [36] T. Pantev and E. Sharpe, *String Compactifications on Calabi–Yau Stacks*, Nucl. Phys. **B733** (2006) 233–296, hep-th/0502044.
- [37] T. Bridgeland, A. King, and M. Reid, *Mukai implies McKay*, J. Amer. Math. Soc. **14** (2001) 535–554, math.AG/9908027.
- [38] T. Bridgeland, *Derived Categories of Coherent Sheaves*, math.AG/0602129, to appear to in proceedings of the ICM 2006.
- [39] P. S. Aspinwall, *D-Branes on Calabi–Yau Manifolds*, in J. M. Maldacena, editor, “Progress in String Theory. TASI 2003 Lecture Notes”, pages 1–152, World Scientific, 2005, hep-th/0403166.

- [40] P. Candelas et al., *Mirror Symmetry for Two Parameter Models — I*, Nucl. Phys. **B416** (1994) 481–562, hep-th/9308083.